

Physiological Differences between Mood Disorder Phenotypes Based on Heart Rate Variability

Marie Casey Olseth^{1,2}, Françoise Crevel^{1,2}, Peta Slocombe^{1,3} and Archie Defillo^{1,2*}

¹Medibio, Minnesota, USA

²West End Medical Consulting, Minnesota, USA

³Vital Conversation Psychology Group, Australia

*Corresponding Author: Archie Defillo, Medibio, West End Medical Consulting, Minnesota, USA.

Received: May 18, 2018

Abstract

Background: Psychiatric conditions have always been diagnosed by clinical interview of patients. Diagnostic accuracy based on this subjective method of psychiatric diagnosis is limited by many factors including skill of the clinician in interviewing psychiatric patients, patient's insight into their symptoms and ability to verbalize distinct characteristics of their symptoms. Use of objective measures for diagnosing psychiatric conditions has not been available to clinicians to improve diagnostic accuracy.

Methods: We performed a retrospective analysis on 24-hour heart rate means and variabilities (standard deviations) of 301 consecutive patients with the diagnosis of either normal, anxiety, depression or mixed. Diagnoses were predicted using multinomial logistic regression analysis. Inter-rater agreement prediction was calculated using Kappa. Mean heart rates of the diagnostic groups were compared simultaneously to the normal cohort using Tukey's range test. Diagnostic accuracy, sensitivity and specificity of the predicted diagnoses compared to the normal cohort were calculated using binomial proportion estimates and ROC curve estimates. Groups were compared to the normal cohort.

Results: Overall Inter-rater agreement was 0.79. Accuracy of diagnosis prediction was 57%. Sensitivities and specificities by diagnoses were: anxiety: 68%, 67%; depression: 35%, 91%; mixed: 0%, 100% (no subjects were diagnosed by the regression as "mixed"). Mean HRs were significantly lower in normal than in the mental illness groups. Standard deviations of the HRs appear to distinguish groups ONLY by the nighttime variation, which is significantly less in the depressed patients than in the normal or anxiety patients. This is especially interesting because it seems to indicate that depressed patients have more consistently higher HRs when compared with anxiety and normal cohort. Equally interesting was the finding of anxiety patients having higher than normal HR, but the same variability as normal patients.

Conclusion: Distinct patterns in mean heart rates and heart rate variability are found that clearly differentiate mood disorder diagnostic groups from normal controls. These reproducible and consistent findings reveal a new opportunity for the improving the accuracy of psychiatric diagnosis through use of these distinct patterns of mean heart rate and heart rate variability.

Keywords: Depression; Anxiety; Heart Rate; Heart Rate Variability

Abbreviations

HR: Heart Rate; HRV: Heart Rate Variability; ROC: Receiver Operating Characteristic

Introduction

Psychiatric conditions have always been diagnosed by clinical interview of patients. Diagnostic accuracy based on this subjective method of psychiatric diagnosis is limited by many factors including skill of the clinician in interviewing psychiatric patients, patient's insight into their symptoms and ability to verbalize distinct characteristics of their symptoms. Use of objective measures for diagnosing psychiatric conditions has not been available to clinicians to improve diagnostic accuracy.

Heart rate and heart rate variability (HRV), specifically time and spectral domains, have been employed in psychiatry conditions (especially depression, anxiety, or both as comorbid), as an accurate, reproducible and non-invasive tool for the assessment of autonomic nervous system dysfunction [1-6]. HRV is now widely investigated as a bio-physiological marker for a variety of mental disorders including, unipolar and bipolar disorders, dysthymia, and schizophrenia [1,5-7].

In mental illness, due to multifactorial reasons, diagnosing particular condition is still challenging by experienced clinicians. The presence of comorbid conditions, such as anxiety and depression, always make the clinical diagnosis more difficult and cumbersome. A unique characteristic of HRV when evaluation and monitoring mental illness is the diversity of patterns that exists between behavioral conditions and the response of HRV to the severity of the symptoms [6,8-13].

Concerning HRV in depression, a reduced parasympathetic activity and a shift of autonomic balance toward sympathetic activation have been demonstrated by researchers in the recent past [6,11,12,14]. The severity of depressive symptoms has been correlated with a profound reduction in parasympathetic modulation [6,11,12,14]. In anxiety disorders, including generalized anxiety disorder, post-traumatic stress disorder, social anxiety disorder, and panic disorder, a reduction in HRV has also been reported [6,16-21]. However, it seems in anxious individual's heart rate changes are more prominent than extreme variations in HRV [21].

Our research demonstrates the value of using autonomic assessment when comparing the patho-physiological profiles of depression and anxiety in order to determine the physiological differences between mood disorder phenotypes.

Materials and Methods

We performed a retrospective analysis on 24-hour heart rate means and variabilities (standard deviations) of 301 consecutive patients with the diagnosis of either normal, anxiety, depression or mixed. Diagnoses were predicted using multinomial logistic regression analysis. Inter-rater agreement prediction was calculated using Kappa. Mean heart rates of the diagnostic groups were compared simultaneously to the normal cohort using Tukey's range test. Diagnostic accuracy, sensitivity and specificity of the predicted diagnoses compared to the normal cohort were calculated using binomial proportion estimates and ROC curve estimates. Groups were compared to the normal cohort.

Results and Discussion

Overall Inter-rater agreement was 0.79. Accuracy of diagnosis prediction was 57%. Sensitivities and specificities by diagnoses were: anxiety: 68%, 67%; depression: 35%, 91%; mixed: 0%, 100% (no subjects were diagnosed by the regression as "mixed"). Mean HRs were significantly lower in normal than in the mental illness groups (Table 1). Standard deviations of the HRs appear to distinguish groups only by the nighttime variation, which is significantly less in the depressed patients than in the normal or anxiety patients. This is especially interesting because it seems to indicate that depressed patients have more consistently higher HRs when compared with anxiety and normal cohort. Equally interesting was the finding of anxiety patients having higher than normal HR, but the same variability as normal patients.

	24hr Heart Rate		Day Heart Rate		Night Heart Rate	
	Mean	SD	Mean	SD	Mean	SD
Condition						
Normal	71.41	7.56	79.14	8.96	63.70	7.49
Anxiety	84.42	9.56	91.81	11.0	77.03	9.35
Depression	83.66	12.7	90.90	14.6	76.42	11.8
Mixed	85.08	10.9	92.34	12.6	77.82	9.9
Groups Compared using Tukey Test						
Anxiety vs Normal	***	NS	***	NS	***	NS
Depression vs Normal	***	NS	***	NS	***	**
Mixed vs Normal	***	NS	***	NS	***	NS
Depression vs Anxiety	NS	NS	NS	NS	NS	**

Table 1: Values of heart rate during the day and the night.

* $P < .05$, ** $P < .01$, *** $P < .001$

The present study intended to examine the physiological differences between mood disorder phenotypes, specifically anxiety, depression or mixed diagnosis and compared to a normal cohort. Our clinical variances were focused on alterations in heart rate and heart rate variability components. The present study examined HR and HRV changes in the early stages of the disease, so we avoided the confounding effects of anti-psychotic or anti-depressive medications. Of notice, diagnosis of anxiety and depression can be challenging for any mental health care provider and when the two disorders coexist diagnostic precision is more complicated.

In our study, we demonstrated that HR and HRV in non-medicated patients were different in the depression group, when compared to anxiety and normal cohort. Depressed patients have more consistently higher HRs when compared with anxiety and normal cohort. Equally interesting was the finding of anxiety patients having higher than normal HR, but the same variability as normal patients. Suggesting that reduction in HRV could be used as a psychophysiological marker between mood disorder phenotypes.

HRV time and spectral domains can be used as intrinsic bio-physiological markers of mental illness [22,23]. These components can accurately, non-invasively, detect cardiac autonomic dysregulation. By means of these factors researchers can identify parasympathetic and sympathetic nervous system over or down modulation in a short term – 24 hours, or over a period of days – longitudinal monitoring [22-32].

It is hypothesized that reduced vagal modulation is accompanied by a subsequent displacement of the sympathovagal balance, this is demonstrated by changes in low frequency/high frequency ratio of HRV- spectral domain. Researchers studied further parameters of HRV- spectral domains such as high frequency power, which estimates parasympathetic activity, and low frequency band which reflects both vagal and sympathetic influences [1,23,33]. Although, Goldstein, *et al.* considers low frequency power to be an index, not of cardiac sympathetic tone, but of baroreflex function during supine rest position [34].

Depressive disorders are associated with sympathetic hyperactivity and reduced vagal control, this is recognized as an independent risk factor of cardiovascular diseases. Similarly, anxiety disorders have been associated with increased risk of cardiovascular morbidity. The latent description for the increased risk of cardiovascular diseases is based on dysregulation of the autonomic nervous system [6,8,11,22].

The cardiovascular signature of depression and anxiety may arise more prominently during sleep stages, due to the sensibility of the heart rate to endogenous dynamic changes across the night like progression of sleep stages, dynamic interactions between homeostatic and circadian processes; these are combined with the fact that sleep minimizes the influence of several external confounders such as fluctuating daytime stress, physical activity and constant cognitive and emotional processing [14,35,36].

Finally, the following clinical theoretical perspectives substantiate our findings. The polyvagal theory initially describes by Porges, it highlights the importance of vagal pathway in attention, emotion expression, social bonding and flexible adjustment to environmental demands. Vagal influences on the heart serve to dampen the sympathetic reaction to stress and to promote calm behavioral states and self-regulation. Without this protective function of vagal tone, which normally disappears or decreases with mental illness, individuals may become vulnerable to anxious apprehension and worry, involving pre-attentive biases to threat information, with rigid and inflexible response patterns, these symptoms are diagnostic traces in patients with anxiety [10,37-45].

Another explanation for decreased parasympathetic tone in anxiety and depression is the inability to disengage threat detection, which serves to perpetuate hyperarousal and worry, even when no real threat exists, which may cause a chronic withdrawal of parasympathetic activity and long-term reduction in HRV [15,37-45].

We understand the limitations of a retrospective analysis, such as: 1) some key statistic factors cannot be measure like control exposure or outcome; 2) we must rely on accurate record-keeping; 3) significant selection biases may affect selection control.

Conclusion

The use of HRV as a transdiagnostic marker has become a novel tool in the psychiatry armamentarium for determination of sympathetic and parasympathetic activity and sympathovagal imbalance. Pathophysiological alterations are likely to alter heart rate dynamics during sleep thereby providing a relevant window to observe multi-systemic characteristic of depression and anxiety. Distinct patterns in mean heart rates and heart rate variability are found that clearly differentiate mood disorder diagnostic groups from normal controls. These reproducible and consistent findings reveal a new opportunity for the improving the accuracy of psychiatric diagnosis through use of these distinct patterns of mean heart rate and heart rate variability.

Acknowledgements

We thank Melissa Martinson, MS PhD for assistance with statistical analysis. All authors had a role in analysis and interpretation of data, in the writing or reviewing of the paper; and in the decision to submit the paper for publication.

Conflict of Interest

Marie Casey Olseth is a part of Medibio's Scientific Advisory Board. The other authors declare that they have no conflicts interests.

Bibliography

1. Alvares GA, et al. "Autonomic nervous system dysfunction in psychiatric disorders and the impact of psychotropic medications: a systemic review and meta-analysis". *Journal of Psychiatry and Neuroscience* 41.2 (2016): 80-104.
2. Berntson GG, et al. "Hear rate variability: origins, methods, and interpretive caveats". *Psychophysiology* 34.6 (1997): 623-648.
3. Thayer JF and Lane RD. "The role of vagal function in the risk for cardiovascular disease and mortality". *Biological Psychology* 74.2 (2007): 224-242.
4. Thayer JF, et al. "The relationship of autonomic imbalance, heart rate variability and cardiovascular disease risk factors". *International Journal of Cardiology* 141.2 (2010): 122-131.
5. Kemp AH, et al. "Impact of depression and antidepressant treatment on heart rate variability: a review and meta-analysis". *Biological Psychiatry* 67.11 (2010): 1067-1074.

Physiological Differences between Mood Disorder Phenotypes Based on Heart Rate Variability

6. Licht CMM., *et al.* "Association between anxiety disorders and heart rate variability in The Netherlands Study of Depression and Anxiety (NESDA)". *Psychosomatic Medicine* 71.5 (2009): 508-518.
7. Quintana DS., *et al.* "A meta-analysis on the impact of alcohol dependence on short term resting-state heart rate variability: implications for cardiovascular risk". *Alcoholism: Clinical and Experimental Research* 37.1 (2013): E23-E29.
8. Gorman JM and Sloan RP. "Heart rate variability in depressive and anxiety disorders". *American Heart Journal* 140.4 (2000): 77-83.
9. Pitzallis MV., *et al.* "Depression but not anxiety influences the autonomic control of heart rate after myocardial infarction". *American Heart Journal* 141.5 (2001): 765-771.
10. Chalmers JA., *et al.* "Worry is associated with robust reductions in heart rate variability: a transdiagnostic study of anxiety psychopathology". *BMC Psychology* 4 (2016): 32.
11. Shinba T. "Major depressive disorder and generalized anxiety disorder show different autonomic dysregulation revealed by heart rate variability analysis in first-onset drug naïve patients without comorbidity". *Psychiatry and Clinical Neurosciences* 71.2 (2017): 135-145.
12. Jahan CR., *et al.* "Autonomic dysfunction in major depressive disorder". *Journal of Bangladesh Society of Physiologists* 9.1 (2014): 37-41.
13. Chang HA., *et al.* "Heart rate variability in unmedicated patients with bipolar disorder in the manic phase". *Psychiatry and Clinical Neurosciences* 68.9 (2014): 674-682.
14. Stampfer HG and Dimmitt SB. "Variations in circadian heart rate in psychiatric disorders: theoretical and practical implications". *Chrono Physiology and Therapy* 3 (2013): 41-50.
15. Hendriks SM., *et al.* "Disorder-specific cognitive profiles in major depressive disorder and generalized anxiety disorder". *BMC Psychiatry* 14 (2014): 96.
16. Harter MC., *et al.* "Association between anxiety disorder and mental illness". *European Archives of Psychiatry and Clinical Neuroscience* 253.6 (2003): 313-320.
17. Vogelzangs N., *et al.* "Cardiovascular disease in person with depressive and anxiety disorder". *Journal of Affective Disorders* 125.1-3 (2010): 241-248.
18. Kemp AH., *et al.* "The association between mood an anxiety disorder, and coronary artery disease in Brazil: a cross-sectional analysis on the Brazilian longitudinal study adult health (ELSA-Brazil)". *Frontiers in Psychology* 6 (2015): 187.
19. Roest AM., *et al.* "Anxiety and risk of incident coronary artery disease: a meta-analysis". *Journal of the American College of Cardiology* 56.1 (2010): 38-46.
20. Janszky I., *et al.* "Early-onset depression anxiety, an risk of subsequent coronary heart disease: 37-year follow-up of 49,321 Swedish men". *Journal of the American College of Cardiology* 56.1 (2010): 31-37.
21. Shibeshi WA., *et al.* "Anxiety worsens prognosis in patients with coronary artery disease". *Journal of the American College of Cardiology* 49.20 (2007): 2021-2027.
22. Wang Y., *et al.* "Altered cardiac autonomic nervous function in depression". *BMC Psychiatry* 13 (2013): 187.
23. Kleiger RE., *et al.* "Heart rate variability: measurement and clinical utility". *Annals of Noninvasive Electrocardiology* 10.1 (2005): 88-101.

Physiological Differences between Mood Disorder Phenotypes Based on Heart Rate Variability

24. Bigger JT, *et al.* "Components of heart rate variability measured during healing of acute myocardial infarction". *American Journal of Cardiology* 61.4 (1988): 208-215.
25. Erwing DJ. "Heart rate variability: and important new risk factor in patients following myocardial infarction". *Clinical Cardiology* 14.8 (1991): 683-685.
26. Kleiger RE., *et al.* "Time domain measurements of heart rate variability". *Cariology Clinics* 10.3 (1992): 487-498.
27. Malik M and Camm J. "Heart rate variability". *Clinical Cardiology* 13 (1990): 570-576.
28. Modanlou HD., *et al.* "A simple method of fetal and neonatal heart rate beat-to-beat variability quantification". *American Journal of Obstetrics and Gynecology* 1279.8 (1997): 861-868.
29. Jennet S., *et al.* "Sudden large and periodic changes in heart rate in healthy young men after short periods of exercise". *British Medical Journal* 285.6349 (1982): 1154-1156.
30. Pagani M., *et al.* "Simultaneous analysis of beat by beat systemic arterial pressure and heart rate variabilities in ambulatory patients". *Journal of Hypertension* 3.3 (1985): S83-S85.
31. Von Neumann J., *et al.* "The mean square successive difference". *Annals of Mathematical Statistics* 12.2 (1941): 153-162.
32. Task force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. "Heart rate variability. Standards of measurements, physiological interpretation, and clinical use". *Circulation* 93.5 (1996): 1043-1065.
33. Quintana DS., *et al.* "Guidelines for reporting articles on psychiatry and heart rate variability (GRAPH): recommendations to advance research communication". *Translational Psychiatry* 6 (2016): e803.
34. Rhaman F., *et al.* "LF power reflects baroreflex function, no cardiac sympathetic innervation". *Clinical Autonomic Research* 21.3 (2011): 133-141.
35. Lanfranchi PA., *et al.* "Cardiac autonomic regulation during sleep in idiopathic REM sleep behavior disorder". *Sleep* 30.8 (2007): 1019-1025.
36. Yan AC., *et al.* "Reduced physiologic complexity is associated with poor sleep in patients with major depression and primary insomnia". *Journal of Affective Disorders* 131.1-3 (2011): 179-185.
37. Porges SW. "The polyvagal theory: neurophysiological foundations of emotions attachment, communication and self-regulation". New York, NY: W.W. Norton and Company (2011).
38. Thayer JF and Lane RD. "A model of neurovisceral integration in emotion regulation and dysregulation". *Journal of Affective Disorders* 61.3 (2000): 201-216.
39. Thayer JF and Lane RD. "Claude Bernard and the heart brain connection: further elaboration of a model of neurovisceral integration". *Neuroscience and Biobehavioral Reviews* 33.2 (2009): 81-88.
40. Clark LA and Watson D. "Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications". *Journal of Abnormal Psychology* 100.3 (1991): 316-336.
41. Clark LA., *et al.* "Temperament, personality, and the mood an anxiety disorder". *Journal of Abnormal Psychology* 103.1 (1994): 103-116.

42. Watson D and Tellegen A. "Toward a consensual structure of mood". *Psychological Bulletin* 98.2 (1985): 219-235.
43. Beck R., *et al.* "The cognitive and emotional phenomenology of depression and anxiety: are worry and hopelessness the cognitive correlates of NA and PA?" *Cognitive Therapy and Research* 25.6 (2001): 829-838.
44. Beck R., *et al.* "Depression and anxiety: integrating the tripartite and cognitive content-specificity models". *Journal of Psychopathology and Behavioral Assessment* 25.4 (2003): 251-256.
45. Jolly JB., *et al.* "Integration of positive and negative affectivity and cognitive content-specificity: improved discrimination of anxious and depressive symptoms". *Journal of Abnormal Psychology* 103.3 (1994): 544-552.

Volume 10 Issue 8 August 2018

©All rights reserved by Eduardo Archie Defillo., *et al.*